

- 35 Standiford TJ, Rolfe MW, Kunkel SL *et al.* Macrophage inflammatory protein-1 alpha expression in interstitial lung disease. *J Immunol* 1993;151:2852–63.
- 36 Bolster MB, Ludwicka A, Sutherland SE *et al.* Cytokine concentrations in bronchoalveolar lavage fluid of patients with systemic sclerosis. *Arthritis Rheum* 1997;40:743–51.
- 37 Hasegawa M, Sato S, Takehara K. Augmented production of chemokines (monocyte chemoattractant protein-1 (MCP-1), macrophage inflammatory protein-1 $\alpha$  (MIP-1 $\alpha$ ) and MIP-1 $\beta$ ) in patients with systemic sclerosis: MCP-1 and MIP-1 $\alpha$  may be involved in the development of pulmonary fibrosis. *Clin Exp Immunol* 1999;117:159–65.
- 38 Zhang Y, McCormick LL, Desai SR *et al.* Murine sclerodermatous graft-versus-host disease, a model for human scleroderma: cutaneous cytokines, chemokines, and immune cell activation. *J Immunol* 2002;168:3088–98.
- 39 Ishida Y, Kimura A, Kondo T *et al.* Essential roles of the CC chemokine ligand 3-CC chemokine receptor 5 axis in bleomycin-induced pulmonary fibrosis through regulation of macrophage and fibrocyte infiltration. *Am J Pathol* 2007;170:843–54.
- 40 Segawa S, Goto D, Yoshiga Y *et al.* Inhibition of transforming growth factor-beta signalling attenuates interleukin (IL)-18 plus IL-2-induced interstitial lung disease in mice. *Clin Exp Immunol* 2010;160:394–402.
- 41 Ihn H. Pathogenesis of fibrosis: role of TGF-beta and CTGF. *Curr Opin Rheumatol* 2002;14:681–5.

## Clinical vignette

Rheumatology 2014;53:2269

doi:10.1093/rheumatology/keu347

Advance Access publication 9 September 2014

### Pneumocephalus due to granulomatosis with polyangiitis

A 67-year-old woman affected by granulomatosis with polyangiitis (GPA) presented with a flare of lung disease and ulceration of the lower limb without active sinusitis. She had a 9-year history of GPA with pulmonary and ENT involvement, the latter leading to left eye enucleation (histological findings suggestive of GPA). She had initially been treated with CYC (6 months) and was currently on steroids and MTX. Laboratory studies revealed raised in-

flammatory markers and cANCA (PR3) positivity. Thoracic CT showed progression of bilateral ground-glass opacities; bronchoalveolar lavage excluded infections. Methylprednisolone 1 mg/kg was introduced with clinical improvement. However, she gradually developed episodic headache and a mild, fluctuant altered mental status with confusion and delayed verbal response without other neurological features. A coronal T2-weighted MRI (Fig. 1) revealed a previously known, unmodified ischaemic lesion and the presence of extensive pneumocephalus secondary to cerebrospinal fluid leakage through multiple discontinuities in the ethmoid bone due to chronic localization of GPA to the anterior skull base (without other risk factors for development of pneumocephalus). Pneumocephalus due to cerebrospinal fluid fistula is usually secondary to trauma or surgery [1], but to our knowledge has never been reported as a complication of skull base involvement in GPA.

**Funding:** None.

**Disclosure statement:** The authors have declared no conflicts of interest.


**Roberto Caporali<sup>1</sup>, Sara Monti<sup>1</sup>, Lorenzo Magrassi<sup>2</sup> and Carlomaurizio Montecucco<sup>1</sup>**

<sup>1</sup>Division of Rheumatology and <sup>2</sup>Division of Neurosurgery, University of Pavia, IRCCS Policlinico San Matteo Foundation, Pavia, Italy.

Correspondence to: Roberto Caporali, Policlinico S. Matteo, Pz. le Golgi 3, 27100 Pavia, Italy.  
E-mail: caporali@smatteo.pv.it

### Reference

- 1 Schirmer CM, Heilman CB, Bhardwaj A. Pneumocephalus: case illustrations and review. *Neurocrit Care* 2010;13:152–8.



**FIG. 1** Coronal T2-weighted MRI revealing the presence of cerebrospinal fluid in the nasal and paranasal cavities (arrow) and air in the left frontal horn of the ventricle (arrowhead)